arXiv:1303.5687v1 [math.AG] 22 Mar 2013

THE GROUP OF UNITS ON AN AFFINE VARIETY

TIMOTHY J. FORD

ABSTRACT. The object of study is the group of units $\mathscr{O}^*(X)$ in the coordinate ring of a normal affine variety X over an algebraically closed field k. Methods of Galois cohomology are applied to those varieties that can be presented as a finite cyclic cover of a rational variety. On a cyclic cover $X \to \mathbb{A}^m$ of affine m-space over k such that the ramification divisor is irreducible and the degree is prime, it is shown that $\mathscr{O}^*(X)$ is equal to k^*, the nonzero scalars. The same conclusion holds, if X is a sufficiently general affine hyperelliptic curve. If X has a projective completion such that the divisor at infinity has r components, then sufficient conditions are given for $\mathscr{O}^*(X)/k^*$ to be isomorphic to $\mathbb{Z}^{(r-1)}$.

1. INTRODUCTION

1.1. **Statement of problem and summary of results.** Throughout, X is a normal variety defined over an algebraically closed field k. If X is an affine variety, and the ring of regular functions on X is denoted $\mathscr{O}(X)$, the group of units $\mathscr{O}^*(X)$ turns out to be a subtle invariant. By comparison, if X is a projective variety in $\mathbb{P}^m$, the ring $\mathscr{O}(X)$ is always equal to k [7, Theorem I.3.4], hence the group of units $\mathscr{O}^*(X)$ is equal to k^*. This confirms the intuitive notion that on a projective variety the only functions with no zero set and no pole set are the non-zero constants.

This article is concerned with the general question, "What is the group of units on an affine variety?" Our main results tend to be one of two types of statements, either sufficient conditions on X such that the group of units is trivial, or sufficient conditions on X such that the group of units is "large". This terminology is explained in Section 1.2. We focus on those varieties X that are either rational, or can be presented as a finite cyclic cover of a rational variety.

In Section 2 the affine variety X is a hypersurface in $\mathbb{A}^{m+1}$ defined by an equation of the form $z^n = f$, where f is a polynomial in $k[x_1, \dots, x_m]$. In Theorems 2.10 and 2.12, we show that $\mathscr{O}^*(X)$ is equal to k^* when f is irreducible and n is either a prime or is equal to 4. The proof relies heavily on Galois cohomology.

In Section 3 the variety X is an affine open subvariety of a finite cyclic covering $\pi : Y \to \mathbb{P}^m$. Section 3.1 treats the case where π is unramified along the divisor at infinity of X. We apply these results to affine hyperelliptic curves in Section 3.2. If $\pi : Y \to \mathbb{P}^1$ is a ramified cyclic cover of prime degree and $k = \mathbb{C}$, then for a sufficiently general $Q \in \mathbb{P}^1$, the group of units on the affine curve $X = \pi^{-1}(\mathbb{P}^1 - Q)$ is trivial (Theorem 3.5). The proofs are based on Galois cohomology. Section 3.3 considers the case where the ramification divisor of π is equal to the divisor at infinity of X. In this case we show that the group of units is large (Proposition 3.12). The proof is based on étale cohomology. These results are applied in Section 3.4 to compute the group of units on an affine variety defined by an equation of the form $f_1 \cdots f_n = 1$, where each f_i is a linear form in $k[x_1, \dots, x_m]$.

Date: March 25, 2013.

2010 *Mathematics Subject Classification.* 13A05; Secondary 14C20, 14R99.

Key words and phrases. affine algebraic variety, group of units, class group.

After stating some familiar background results, Section 1 concludes with examples meant to provide motivation for the rest of the article.

1.2. **Background results.** For background definitions and theorems on group cohomology, we refer the reader to [11]. For étale cohomology, we suggest [8], and for all other unexplained terminology and notation, [7].

Lemma 1.1 is from [12, Lemme 1] and reduces the computation of $\mathscr{O}^*(X)$ as an abstract group to the determination of the rank of a free $\mathbb{Z}$-module. Given the affine variety X and a projective completion Y, we see that the group of units on X is closely tied to the components of the divisor at infinity and the subgroup that they generate in the class group $\mathrm{Cl}(Y)$. The rank of the free $\mathbb{Z}$-module $\mathscr{O}^*(X)/k^*$ is bounded above by r, the number of irreducible components of the divisor at infinity. The group $\mathscr{O}^*(X)$ is trivial, if it is equal to k^*. The group $\mathscr{O}^*(X)$ is said to be *large*, if the rank of $\mathscr{O}^*(X)/k^*$ is greater than or equal to $r-1$. We sketch Samuel's proof. Lemma 1.2 shows that the affine cone over a projective variety has trivial group of units. The proof can be found in [3]. In Lemma 1.3 we compute the group of units on a basic Zariski open affine subset of $\mathbb{A}^m$.

Lemma 1.1. *If X is a normal affine variety, the group $\mathscr{O}^*(X)/k^*$ is a finitely generated torsion free $\mathbb{Z}$-module.*

Proof. Embed X as an open affine subvariety of a normal projective variety Y. Write the "divisor at infinity" as a union of prime divisors $Y-X = Z_1 \cup \cdots \cup Z_r$. As previously remarked, $\mathscr{O}^*(Y) = k^*$. The claim follows from the following version of the exact sequence

$$(1) \qquad 1 \to \mathscr{O}^*(Y) \to \mathscr{O}^*(X) \xrightarrow{\mathrm{div}} \bigoplus_{i=1}^{r} \mathbb{Z}Z_i \to \mathrm{Cl}(Y) \to \mathrm{Cl}(X) \to 0$$

of Nagata (for example, [3, Theorem 1.1]). □

Lemma 1.2. *Let $f \in k[x_0,\dots,x_m]$ be a square-free homogeneous polynomial of degree $d>0$, defining a projective hypersurface V in $\mathbb{P}^m$. Let $X = Z(f)$ be the affine cone over V in $\mathbb{A}^{m+1}$. Then $\mathscr{O}^*(X) = k^*$.*

Lemma 1.3. *Let $A = k[x_1,\dots,x_m]$ be the coordinate ring of $\mathbb{A}^m$. Let $f \in A$ be non-invertible and square-free with factorization $f = f_1 \cdots f_v$ into irreducible elements of A. If $R = A[f^{-1}]$, then*
(a) $A^* = k[x_1,\dots,x_m]^* = k^*$.
(b) $R^* = k[x_1,\dots,x_m][f^{-1}]^* = k^* \times \langle f_1 \rangle \times \cdots \times \langle f_v \rangle$.

Proof. The factorization of f is unique because A is a unique factorization domain. In addition, $\mathrm{Cl}(A) = \mathrm{Cl}(R) = \langle 0 \rangle$. The counterpart of (1) for the localization $R = A[f^{-1}]$ is

$$(2) \qquad 1 \to A^* \to R^* \xrightarrow{\mathrm{div}} \bigoplus_{i=1}^{v} \mathbb{Z}F_i \to \mathrm{Cl}(A) \to \mathrm{Cl}(R) \to 0$$

where $F_i = Z(f_i)$, for $i = 1,\dots,f_v$. The sequence splits because $\bigoplus \mathbb{Z}F_i$ is free. □

1.3. **Motivational examples.**

Example 1.4. Let $m>1$ and $n>1$. Let $A = k[x_1,\dots,x_m]$ be the affine coordinate ring for $\mathbb{A}^m$. In this example we construct an affine variety X of degree n such that the group $\mathscr{O}^*(X)/k^*$ has rank $n-1$. Let $f = f_1 f_2 \dots f_n + 1$ be an irreducible element of A where $f_1,\dots,f_n$ are linear polynomials. In $\mathbb{A}^m$, let $X = Z(f)$, $F_1 = Z(f_1),\dots,F_n = Z(f_n)$ be the affine varieties. Write $(\,)^*$ for homogenization with respect to the variable x_0. Let $\bar{X} =$

$Z(f^*)$, $\bar{F}_1 = Z(f_1^*), \dots, \bar{F}_n = Z(f_n^*)$ be the projective completions of the affine varieties. Let $F_0 = Z(x_0)$ denote the hyperplane at infinity. If we set $L_i = F_i \cdot F_0$, then L_i is a hyperplane in F_0. Assume the polynomials $f_1, \dots, f_n$ are chosen so that X is normal and $L_1, \dots, L_n$ are distinct. In this notation, sequence (1) becomes

$$1 \to k^* \to \mathscr{O}^*(X) \xrightarrow{\text{div}} \bigoplus_{i=1}^{n} \mathbb{Z}L_i \xrightarrow{\chi} \text{Cl}(\bar{X}) \to \text{Cl}(X) \to 0 \tag{3}$$

Let H denote the image of χ, which is the subgroup of $\text{Cl}(\bar{X})$ generated by the divisors $L_1, \dots, L_n$. We will prove the following.

(a) H is an infinite group.
(b) H is a homomorphic image of $\mathbb{Z} \oplus (\mathbb{Z}/n)^{(n-2)}$.
(c) The image of div has rank $n-1$.
(d) The elements $f_1, \dots, f_{n-1}$ generate a subgroup of finite index in $\mathscr{O}^*(X)/k^*$.

The free group $\mathbb{Z}L_i$ maps onto the subgroup of $\text{Cl}(\bar{X})$ generated by L_i. Since L_i has degree one, the degree map $D \mapsto \deg(D)L_i$ is a splitting map. Therefore (a) is true. This also shows the image of div has rank at most $n-1$. At the generic point of $\bar{X}$ we have

$$\frac{f_1^*}{x_0} \cdots \frac{f_n^*}{x_0} = 1$$

so the function f_i^*/x_0 represents a unit in the coordinate ring $\mathscr{O}(X)$. One can compute the intersection divisors $X \cdot F_1 = nL_1, \dots, X \cdot F_n = nL_n, X \cdot F_0 = L_1 + \cdots + L_n$ on $\bar{X}$. Therefore, the divisor of f_i^*/x_0 on $\bar{X}$ is

$$\text{div}(f_i^*/x_0) = nL_i - L_1 - \cdots - L_n. \tag{4}$$

Using (4) one checks that the image of χ is generated by the classes of the $n-1$ divisors $L_n, L_1 - L_n, \dots, L_{n-2} - L_n$. It follows from (4) that

$$\begin{aligned} \text{div}(f_i^*/f_j) &= (nL_i - L_1 - \cdots - L_n) - (nL_j - L_1 - \cdots - L_n) \\ &= n(L_i - L_j). \end{aligned} \tag{5}$$

Equations (4) and (5) can be used to prove (c) and (d). Equation (5) shows that $L_1 - L_n, \dots, L_{n-2} - L_n$ generate a subgroup of $\text{Cl}(\bar{X})$ which is annihilated by n. Therefore (b) is true. We have shown that $\mathscr{O}^*(X)/k^* \cong (\mathbb{Z})^{n-1}$ and the elements $f_1, \dots, f_{n-1}$ generate a subgroup of finite index. Do the elements $f_1, \dots, f_{n-1}$ generate the group $\mathscr{O}^*(X)/k^*$? For some particular cases, Examples 1.5 and 1.6, prove that the answer is yes. Another affirmative answer is given in Example 3.14, where we impose the additional hypotheses on X that $\bar{X} \to \mathbb{P}^m$ is cyclic, and the ramification divisor is $\bar{X} - X$, the divisor at infinity.

Example 1.5. In Example 1.4, if $n \le m$, and $f = x_1 \cdots x_n + 1$, then in $\mathscr{O}(X)$ we have $x_n = -x_1^{-1} \cdots x_{n-1}^{-1}$. The isomorphism

$$\mathscr{O}(X) = \frac{k[x_1, \dots, x_m]}{(x_1 \cdots x_n + 1)} \cong k[x_1, \dots, x_m][x_1^{-1}, \dots, x_{n-1}^{-1}]$$

is defined by eliminating x_n. By Lemma 1.3, $\mathscr{O}^*(X)$ is equal to $k^* \times \langle f_1 \rangle \times \cdots \times \langle f_{n-1} \rangle$.

Example 1.6. In Example 1.4, if $n = 3$, the image of χ is generated by L_3 and $L_1 - L_3$. One can check that $L_1 - L_3$ generates a subgroup of order 3 in $\text{Cl}(\bar{X})$. Therefore the image of χ is equal to the internal direct sum $\mathbb{Z}L_3 \oplus \mathbb{Z}/3(L_1 - L_3)$ and the group $\mathscr{O}^*(X)$ is equal to $k^* \times \langle f_1 \rangle \times \langle f_2 \rangle$.

Example 1.7. A class of curves that have been widely studied are the affine Fermat curves $F = Z(x^n + y^n - 1)$ in $\mathbb{A}^2$, where $n \geq 2$. The argument in Example 1.4 shows that the group $\mathscr{O}^*(F)/k^*$ is free of rank $n-1$. Let $\bar{F}$ denote the projective completion, and X the affine open where $xy \neq 0$. Let H denote the subgroup of $\mathrm{Cl}(\bar{F})$ generated by the $3n$ points in $\bar{F} - X$. If H_0 is the kernel of the degree homomorphism, then

$$0 \to H_0 \to H \xrightarrow{\deg} \mathbb{Z} \to 0 \tag{6}$$

is a split exact sequence, because a point has degree one. As shown in [13], H_0 is a finite group which is annihilated by n. Using Lemma 1.1 we find that $\mathscr{O}^*(X)/k^*$ has rank $3n-1$. A basis for $\mathscr{O}^*(X)/k^*$ is computed in [10].

2. A CYCLIC COVER OF AFFINE SPACE

In this section we consider an affine variety X which is a finite cyclic cover of $\mathbb{A}^m$. It will be helpful to set up some notation. Let $A = k[x_1, \dots, x_m]$ be the coordinate ring of $\mathbb{A}^m$. Let f be a non-invertible square-free element of A. Assume $n \geq 2$ is invertible in k and ζ is a primitive nth root of unity in k. Set $T = A[z]/(z^n - f)$ be $\mathscr{O}(X)$, the coordinate ring of the affine variety X. Then T is a ramified cyclic extension of A, and the associated morphism on varieties is $\pi : X \to \mathbb{A}^m$. Let $f = f_1 \dots f_v$ be a factorization of f into irreducibles in A. By Eisenstein's criterion, for instance, T is an integral domain. The singular locus of X corresponds to the singular locus of the affine hypersurface $F = Z(f) \subseteq \mathbb{A}^m$. Hence X is regular in codimension one. By the Serre criteria, X is normal, and T is an integrally closed integral domain. The assignment $\sigma(z) = \zeta z$ defines an A-algebra automorphism of T. If $G = \langle \sigma \rangle$, then $T^G = A$. The morphism π ramifies only above the divisor F. If we let $R = A[f^{-1}]$ and $S = T[z^{-1}]$, then S is a Galois extension of R with cyclic group G. The rings defined so far make up this commutative diagram

$$\begin{tikzcd} T = A[\sqrt[n]{f}] \arrow[r] & S = R[\sqrt[n]{f}] \\ A \arrow[u] \arrow[r] & R = A[f^{-1}] \arrow[u] \end{tikzcd} \tag{7}$$

where an arrow represents set inclusion. The norm

$$N : T^* \to k^* \tag{8}$$

is a homomorphism of abelian groups, defined by $N(u) = u\sigma(u)\cdots\sigma^{n-1}(u)$. If $a \in k^*$, then $N(a) = a^n$ because k is a subfield of A. Since k is algebraically closed, N is onto.

2.1. The group of units on a cyclic cover of affine space.

The notation established in the preceding paragraph is used throughout this section.

Proposition 2.1. *In the above context, the following are true.*

(a) $$\mathrm{H}^i(G, k^*) = \begin{cases} k^* & \text{if } i = 0 \\ \mu_n & \text{if } i = 1, 3, 5, \dots \\ \langle 1 \rangle & \text{if } i = 2, 4, 6, \dots \end{cases}$$

(b) $$\mathrm{H}^i(G, R^*) = \begin{cases} R^* = k^* \times \langle f_1 \rangle \times \cdots \times \langle f_v \rangle & \text{if } i = 0 \\ \mu_n & \text{if } i = 1, 3, 5, \dots \\ \frac{\langle f_1 \rangle}{\langle f_1^n \rangle} \times \cdots \times \frac{\langle f_v \rangle}{\langle f_v^n \rangle} & \text{if } i = 2, 4, 6, \dots \end{cases}$$

(c) $$\mathrm{H}^i(G,R^*/k^*) = \begin{cases} R^*/k^* = \langle f_1\rangle \times \cdots \times \langle f_v\rangle & \text{if } i=0 \\ \langle 1\rangle & \text{if } i=1,3,5,\ldots \\ \frac{\langle f_1\rangle}{\langle f_1^n\rangle} \times \cdots \times \frac{\langle f_v\rangle}{\langle f_v^n\rangle} & \text{if } i=2,4,6,\ldots \end{cases}$$

Proof. The sequence of trivial G-modules

$$1 \to k^* \to R^* \to R^*/k^* \to 1 \tag{9}$$

is exact. Use Lemma 1.3 and [11, Theorem 10.35]. □

Lemma 2.2. *In the context of Section 2.1, let H be a subgroup of G of order h. If $(T^*)^H = k^*$, then the following are true.*

(a) $(T^*/k^*)^H = \langle 1\rangle$.

(b) $\mathrm{H}^{2i}(H,T^*/k^*) = \langle 1\rangle$, *for all* $i > 0$.

(c) There is an exact sequence

$$1 \to \mu_h \to \mathrm{H}^{2i-1}(H,T^*) \to \mathrm{H}^{2i-1}(H,T^*/k^*) \to 1$$

for all $i > 0$.

Proof. Begin with the exact sequence

$$1 \to k^* \to T^* \to T^*/k^* \to 1. \tag{10}$$

By Lemma 1.1, T^*/k^* is a finitely generated torsion free $\mathbb{Z}$-module. The long exact sequence associated to (10) is

$$\begin{aligned} &1 \to k^* \xrightarrow{\alpha^0} (T^*)^H \to (T^*/k^*)^H \xrightarrow{\partial^0} \\ &\quad \mathrm{H}^1(H,k^*) \xrightarrow{\alpha^1} \mathrm{H}^1(H,T^*) \to \mathrm{H}^1(H,T^*/k^*) \xrightarrow{\partial^1} \\ &\quad\quad \mathrm{H}^2(H,k^*) \xrightarrow{\alpha^2} \mathrm{H}^2(H,T^*) \to \mathrm{H}^2(H,T^*/k^*) \xrightarrow{\partial^2} \ldots \end{aligned} \tag{11}$$

By hypothesis, α^0 is an isomorphism, so ∂^0 is one-to-one. Since G acts trivially on k^*, $\mathrm{H}^1(H,k^*) = \mu_h$, a finite group. Because $(T^*/k^*)^H$ is a subgroup of the torsion free group T^*/k^*, we conclude that (a) is true. Because H is a cyclic group, part (b) follows from [11, Theorem 10.35] and part (a). By Proposition 2.1(a), for $i > 0$, $\mathrm{H}^{2i}(H,k^*) = \langle 1\rangle$. Part (c) follows from (11), part (a) and periodicity. □

Theorem 2.3. *In the context of Section 2.1, assume $n = p$ is a prime number. There is an isomorphism of $\mathbb{Z}[G]$-modules $T^*/k^* \cong A_1 \oplus \cdots \oplus A_t$ where $A_1,\ldots,A_t$ are $\mathbb{Z}[\zeta]$-ideals in $\mathbb{Q}[\zeta]$. It follows that T^*/k^* is a free $\mathbb{Z}$-module of rank $(p-1)t$ and there are isomorphisms*

$$\mathrm{H}^1(G,T^*) \cong \mu_p \times (\mathbb{Z}/p)^{(t)}$$
$$\mathrm{H}^1(G,T^*/k^*) \cong (\mathbb{Z}/p)^{(t)}$$

Proof. Since $(T^*)^G = k^*$, this follows from Lemma 2.2 and the structure theory for a module over a cyclic group of order p, [2, Theorem (74.3)]. A similar argument is given in [4, Proposition 2.17]. □

Example 2.4. When $v > 1$, the group of units of T can be trivial. For instance, let $f = (xy-1)(x-\alpha_1)\cdots(x-\alpha_d)$, where $\alpha_1,\ldots,\alpha_d$ are distinct elements in k^*. As computed in [5, §3.3], the group of units in $T = A[z]/(z^2-f)$ is k^*.

The group G acts as a group of automorphisms of $\mathrm{Cl}(T)$. This action is induced on the group of divisors $\mathrm{Div}(T)$ by sending a height one prime ideal I to its conjugate $\sigma(I)$. By this same action, G acts as a group of automorphisms of $\mathrm{Pic}(S)$. Because $\operatorname{Spec} S$ is nonsingular, $\mathrm{Pic}\,(S) = \mathrm{Cl}\,(S)$. Associated to the Galois extension S/R is

$$\begin{aligned} (12)\quad & 1 \to \mathrm{H}^1(G,S^*) \xrightarrow{\alpha_1} \mathrm{Pic}(R) \xrightarrow{\alpha_2} (\operatorname{Pic} S)^G \xrightarrow{\alpha_3} \\ & \mathrm{H}^2(G,S^*) \xrightarrow{\alpha_4} \mathrm{B}(S/R) \xrightarrow{\alpha_5} \mathrm{H}^1(G,\operatorname{Pic} S) \xrightarrow{\alpha_6} \mathrm{H}^3(G,S^*) \end{aligned}$$

the so-called Chase-Harrison-Rosenberg exact sequence of cohomology groups [1, Corollary 5.5]. The group $\mathrm{B}(S/R)$ appearing in (12) is the kernel of the natural map on Brauer groups $\mathrm{B}(R) \to \mathrm{B}(S)$. In this article it plays only a peripheral role. Since $\operatorname{Pic} R = 0$, sequence (12) and periodicity implies

$$\mathrm{H}^i(G,S^*) = \langle 1 \rangle \tag{13}$$

if $i > 0$ is odd.

Proposition 2.5. *In the context of Section 2.1, the following are true.*

(a) If $i \geq 0$ is even, then

$$\mathrm{H}^i(G,S^*/R^*) = (S^*/R^*)^G \cong \mathbb{Z}/n$$

is generated by the coset containing z.

(b) If $i > 0$ is odd, there is an exact sequence

$$1 \to \mathrm{H}^i(G,S^*/R^*) \to \mathrm{H}^{i+1}(G,R^*) \to \mathrm{H}^{i+1}(G,S^*) \to 1$$

of $\mathbb{Z}/n$-modules.

Proof. The exact sequence of G-modules

$$1 \to R^* \to S^* \to S^*/R^* \to 1 \tag{14}$$

gives rise to the exact sequence of cohomology

$$\begin{aligned} (15)\quad & 1 \to R^* \to (S^*)^G \to (S^*/R^*)^G \xrightarrow{\partial^0} \mathrm{H}^1(G,R^*) \to \mathrm{H}^1(G,S^*) \to \\ & \mathrm{H}^1(G,S^*/R^*) \xrightarrow{\partial^1} \mathrm{H}^2(G,R^*) \to \mathrm{H}^2(G,S^*) \to \\ & \mathrm{H}^2(G,S^*/R^*) \xrightarrow{\partial^2} \mathrm{H}^3(G,R^*) \to \mathrm{H}^3(G,S^*) \to \dots \end{aligned}$$

Equation (13) implies ∂^0 and ∂^2 are onto. Since $R^* = (S^*)^G$, ∂^0 is an isomorphism. The minimum polynomial of z is $z^n - f$. In S^*/R^*, the coset containing z has order n. The image of the norm map $N : S^*/R^* \to S^*/R^*$ is $\langle 1 \rangle$. This proves (a). Since ∂^2 is also an isomorphism, we get (b). ☐

Proposition 2.6. *In the context of Section 2.1, the following are true*

(a) $(S^*/k^*)^G = \langle z \rangle \times \langle f_2 \rangle \times \cdots \times \langle f_\nu \rangle$ is a free $\mathbb{Z}$-module of rank ν.

(b) If $i > 0$ is odd, then

$$\mathrm{H}^i(G,S^*/k^*) = \langle 1 \rangle.$$

(c) If $i > 0$ is even, there is an exact sequence

$$1 \to \mathrm{H}^i(G,S^*) \to \mathrm{H}^i(G,S^*/k^*) \to \mathbb{Z}/n \to 1$$

of $\mathbb{Z}/n$-modules.

Proof. In the exact sequence of G-modules

$$1 \to R^*/k^* \to S^*/k^* \xrightarrow{\eta} S^*/R^* \to 1 \tag{16}$$

$\eta(z) = z$. The sequence of cohomology associated to (16) is

$$1 \to R^*/k^* \to (S^*/k^*)^G \xrightarrow{\eta} (S^*/R^*)^G \xrightarrow{\partial^0} \mathrm{H}^1(G, R^*/k^*) \to \dots. \tag{17}$$

By Proposition 2.1, in (17), η is onto. By Proposition 2.5, the image of η is generated by $\eta(z)$. So $(S^*/R^*)^G$ is an extension of R^*/k^* by the finite cyclic group $\langle z\rangle$. By Lemma 1.1, $(S^*/k^*)^G$ is a finitely generated torsion free $\mathbb{Z}$-module. By Lemma 1.3, $R^*/k^* = \langle f_1\rangle \times \cdots \times \langle f_v\rangle$. Since $z^n = f_1 \cdots f_v$, this implies $(S^*/k^*)^G$ is generated by $z, f_2, \dots, f_v$, proving (a).

The exact sequence of G-modules

$$1 \to k^* \to S^* \to S^*/k^* \to 1 \tag{18}$$

gives the exact sequence of cohomology

$$\begin{aligned} &\mathrm{H}^{2i-1}(G,S^*) \to \mathrm{H}^{2i-1}(G,S^*/k^*) \xrightarrow{\partial^{2i-1}} \mathrm{H}^{2i}(G,k^*) \\ &\to \mathrm{H}^{2i}(G,S^*) \to \mathrm{H}^{2i}(G,S^*/k^*) \xrightarrow{\partial^{2i}} \mathrm{H}^{2i+1}(G,k^*) \to \mathrm{H}^{2i+1}(G,S^*) \to \end{aligned} \tag{19}$$

where $i > 0$ is arbitrary. Parts (b) and (c) follow from (19), (13), and Proposition 2.1(a). □

Example 2.7. When $v > 1$, the group of units of T can be non-trivial. For instance, let $f = (xy-1)(xy+1) = f_1 f_2$. As computed in [5, § 3.2], the group of units in $T = A[z]/(z^2 - f)$ is

$$T^* = k^* \times \langle z - xy\rangle \tag{20}$$

and the group of units in $S = T[z^{-1}]$ is

$$S^* = k^* \times \langle z-xy\rangle \times \langle z-xy+1\rangle \times \langle z-xy-1\rangle. \tag{21}$$

Use the identity $\sigma(z-xy) = (z-xy)^{-1}$ to compute

$$\mathrm{H}^i(G,T^*) = \begin{cases} k^* & \text{if } i = 0 \\ \mu_2 \times \frac{\langle z-xy\rangle}{\langle (z-xy)^2\rangle} & \text{if } i = 1,3,5,\dots \\ \langle 1\rangle & \text{if } i = 2,4,6,\dots \end{cases} \tag{22}$$

and

$$\mathrm{H}^i(G,T^*/k^*) = \begin{cases} \langle 1\rangle & \text{if } i \text{ is even} \\ \frac{\langle z-xy\rangle}{\langle (z-xy)^2\rangle} & \text{if } i \text{ is odd.} \end{cases} \tag{23}$$

Use the identities

$$\begin{aligned} \sigma(z-xy+1) &= (z-xy+1)(z-xy)^{-1} \\ \sigma(z-xy-1) &= -(z-xy-1)(z-xy)^{-1} \\ 2z &= (z-xy+1)(z-xy-1)(z-xy)^{-1} \\ -2(xy-1) &= (z-xy+1)^2(z-xy)^{-1} \\ -2(xy+1) &= (z-xy-1)^2(z-xy)^{-1} \end{aligned}$$

to compute

$$\mathrm{H}^i(G,S^*) = \begin{cases} R^* = k^* \times \langle xy-1\rangle \times \langle xy+1\rangle & \text{if } i=0 \\ \langle 1\rangle & \text{if } i>0 \end{cases} \tag{24}$$

and

$$\mathrm{H}^i(G,S^*/k^*) = \begin{cases} \langle z\rangle \times \langle xy+1\rangle & \text{if } i=0 \\ \langle 1\rangle & \text{if } i=1,3,5,\dots \\ \frac{\langle z\rangle}{\langle z^2\rangle} & \text{if } i=2,4,6,\dots. \end{cases} \tag{25}$$

These results agree with Propositions 2.5 and 2.6.

2.2. **The ramification divisor is irreducible.** In addition to the notation established in the opening paragraph of Section 2, assume that $T = A[z]/(z^n - f)$, where f is irreducible in A.

Theorem 2.8. *In the context above, the following are true.*
(a) $\mathrm{Cl}(T) = \mathrm{Cl}(S)$.
(b) $\mathrm{Cl}(T)^G = \mathrm{Cl}(S)^G = \langle 0\rangle$.
(c)

$$\mathrm{H}^i(G,T^*) = \begin{cases} k^* & \textit{if } i=0 \\ \cong \mathbb{Z}/n & \textit{if } i=1,3,5,\dots \\ \langle 1\rangle & \textit{if } i=2,4,6,\dots \end{cases}$$

(d)

$$\mathrm{H}^i(G,S^*) = \begin{cases} R^* = k^* \times \langle f\rangle & \textit{if } i=0 \\ \langle 1\rangle & \textit{if } i>0 \end{cases}$$

(e) $\mathrm{B}(S/R) \cong \mathrm{H}^1(G, \mathrm{Pic}\, S)$.

Proof. Since $T/(z) = A/(f)$, the ideal $I = Tz$ is a height one prime. In T, $z^n = f$, so the ideal Tf has only one minimal prime, namely the height one prime $I = Tz$. By Nagata's Theorem [6, Theorem 7.1], the sequence

$$1 \to T^* \to S^* \xrightarrow{\mathrm{div}} \mathbb{Z}I \to \mathrm{Cl}(T) \to \mathrm{Cl}(S) \to 0 \tag{26}$$

is exact. The divisor of z is $\mathrm{div}(z) = I$, so (26) shows $\mathrm{Cl}(T) = \mathrm{Cl}(S)$. It also follows from (26) that $S^*/T^* = \langle z\rangle$. Since $\sigma(z) = \zeta z$ and $\zeta \in T^*$, we see that S^*/T^* is a trivial G-module. Therefore,

$$\mathrm{H}^i(G,S^*/T^*) = \begin{cases} \langle z\rangle & \text{if } i=0, \\ 1 & \text{if } i=1,3,5,\dots, \\ \langle z\rangle/\langle z^n\rangle & \text{if } i=2,4,6,\dots \end{cases} \tag{27}$$

[11, Theorem 10.35]. The long exact sequence associated to $1 \to T^* \to S^* \to S^*/T^* \to 1$ is

$$1 \to (T^*)^G \to (S^*)^G \to S^*/T^* \to \mathrm{H}^1(G,T^*) \to \mathrm{H}^1(G,S^*) \to \mathrm{H}^1(G,S^*/T^*) \to \dots \tag{28}$$

As in Lemma 1.3, $\mathrm{Pic}(R) = \langle 0\rangle$. By sequence (12), $\mathrm{H}^i(G,S^*) = \langle 1\rangle$ for odd i. Again by Lemma 1.3, $(T^*)^G = A^* = k^*$. The norm map (8) is onto, so for $i = 2,4,6,\dots$, we have $\mathrm{H}^i(G,T^*) = 1$. By Lemma 1.3, $(S^*)^G = R^* = k^* \times \langle f\rangle$. The terms of lowest degree in (28) give rise to the short exact sequence

$$1 \to \langle f\rangle \to \langle z\rangle \to \mathrm{H}^1(G,T^*) \to 1. \tag{29}$$

Therefore $\mathrm{H}^i(G,T^*) = \langle z\rangle/\langle z^n\rangle \cong \mathbb{Z}/n$ for odd i, proving (c). On the element z, the norm map $N : S^* \to R^*$ is

$$N(z) = z\zeta z\cdots\zeta^{n-1}z = \zeta^{n(n-1)/2}f. \tag{30}$$

Use this and the fact that (8) is onto to prove that $\mathrm{H}^i(G,S^*) = 1$ for $i = 2,4,6,\dots$, proving (d). Part (b) follows from (a), (d), and sequence (12). Part (e) follows from sequence (12). ☐

Conjecture 2.9. *If f is irreducible, then $k^* = T^*$. A partial answer is given in Theorems 2.10 and 2.12.*

Theorem 2.10. *In the context of Section 2.2, assume $n = p$ is a prime number. Then $k^* = T^*$.*

Proof. This is a consequence of Theorem 2.3 and Theorem 2.8(c). ☐

Lemma 2.11. *Let p be a prime and $n = p^2$. Let $G = \langle\sigma\rangle$, $H = \langle\sigma^p\rangle$. Let*

$$\begin{aligned} T_2 &= A[z]/(z^n - f),\\ S_2 &= R[z]/(z^n - f),\\ T_1 &= T_2^H = A[z^p],\\ S_1 &= S_2^H = R[z^p]. \end{aligned}$$

The following are true.

(a)

$$\mathrm{H}^i(H,S_2^*/T_2^*) = \begin{cases} \langle z\rangle & \text{if } i = 0\\ \langle 1\rangle & \text{if } i = 1,3,5,\dots\\ \langle z\rangle/\langle z^p\rangle & \text{if } i = 2,4,6,\dots. \end{cases}$$

(b) $\mathrm{H}^0(H,T_2^*) = T_1^* = k^*$ *and* $\mathrm{H}^0(H,S_2^*) = S_1^* = k^* \times \langle z^p\rangle$.

(c) If $i > 0$ is even, then $\mathrm{H}^i(H,T_2^*) = 1$ *and* $\mathrm{H}^i(H,S_2^*) = 1$.

(d) If $i > 0$ is odd, there is a short exact sequence

$$1 \to \mu_p \to \mathrm{H}^i(H,T_2^*) \to \mathrm{H}^i(H,S_2^*) \to 1.$$

(e) The group $\mathrm{H}^1(H,S_2^*)$ *is isomorphic to* $\mathrm{Cl}(S_2/S_1)$*, which is the kernel of the natural map* $i : \mathrm{Cl}(S_1) \to \mathrm{Cl}(S_2)$.

Proof. From Theorem 2.8, $S_2^*/T_2^* = \langle z\rangle$ with trivial G-action. This gives (a). Since $T_2^H = T_1$, Theorem 2.10 implies $(T_2^*)^H = T_1^* = k^*$ and $(S_2^*)^H = S_1^* = k^* \times \langle z^p\rangle$, which is (b). Since H is cyclic, if $i > 0$ is even, then $\mathrm{H}^i(H,T_2^*)$ is equal to k^*/NT_2^*, which is trivial for the same reason that the norm (8) is onto. As in (30), one can check that z^p is in the image of the norm map $S_2^* \to S_1^*$. Part (c) follows. The exact sequence

$$\begin{aligned} &1 \to (T_2^*)^H \to (S_2^*)^H \to S_2^*/T_2^* \to \mathrm{H}^1(G,T_2^*) \to\\ &\qquad \mathrm{H}^1(H,S_2^*) \to \mathrm{H}^1(H,S_2^*/T_2^*) \to \mathrm{H}^2(H,T_2^*) \to \mathrm{H}^2(H,S_2^*) \to\\ &\qquad\quad \mathrm{H}^2(H,S_2^*/T_2^*) \to \mathrm{H}^3(H,T_2^*) \to \mathrm{H}^3(H,S_2^*) \to \mathrm{H}^3(H,S_2^*/T_2^*) \to \dots \end{aligned} \tag{31}$$

is the counterpart of (28) for the group H. Use (31), parts (a), (b), (c), and periodicity to get (d). The groups in (d) are $\mathbb{Z}/p$-modules, so the sequence splits. Part (e) is proved in [6, Theorem 16.1]. ☐

Theorem 2.12. *If $n = 4$, then $k^* = T^*$.*

Proof. In Lemma 2.11(e), the group $\mathrm{Cl}(S_2/S_1)$ is a subgroup of ${}_2\mathrm{Cl}(S_1)$. By [5, Theorem 2.3], ${}_2\mathrm{Cl}(S_1)$ is equal to $\mathrm{Cl}(S_1)^{G/H}$, which by Theorem 2.8(b) is equal to $\langle 0\rangle$. Lemma 2.11(e) implies $\mathrm{H}^1(H,S_2^*) = \langle 1\rangle$ and Lemma 2.11(d) implies $\mathrm{H}^1(H,T_2^*) \cong \mu_2$. By Lemma 2.2, $\mathrm{H}^1(H,T_2^*/k^*) = \langle 1\rangle$. As in the proof of Theorem 2.10, $T_2^*/k^* = \langle 1\rangle$. □

Conjecture 2.13. *If $n = 2^s$, for any $s > 0$, then $k^* = T^*$. To iterate the argument used in Theorem 2.12, it is necessary to know that elements of order two in $\mathrm{Cl}(S_i)$ are fixed by G.*

3. LOCALIZATION OF A CYCLIC COVER OF PROJECTIVE SPACE

In this section we consider the group of units on a ramified cyclic cover $\pi : X \to U$ of affine varieties which is the restriction of a cyclic cover of projective space $\pi : Y \to \mathbb{P}^m$ to an open set. We treat two special cases. In Section 3.1, the localization of π is such that along the “divisor at infinity”, π is unramified. Section 3.3 considers the case where the “divisor at infinity” is equal to the ramification divisor. In Section 3.2, these results are applied to the group of units on an affine curve. In Section 3.4, the results of this section are applied to a special case of Example 1.4.

3.1. Unramified at infinity.

Start with a normal projective variety Y of dimension $m > 0$, together with $G = \langle\sigma\rangle$ a cyclic group of order n acting on Y such that the quotient morphism is $\pi : Y \to \mathbb{P}^m$. Next, we restrict π to an affine open subset of $\mathbb{P}^m$ which is the complement of a prime divisor that is split by π. Specifically, let $F \subseteq \mathbb{P}^m$ be an irreducible hypersurface and $U = \mathbb{P}^m - F$ the affine open complement. Assume the irreducible hypersurface $F \subseteq \mathbb{P}^m$ is split by π. By this we mean $\pi : Y \times_{\mathbb{P}^m} F \to F$ is unramified, and lying above F are n irreducible components. If we write $\pi^{-1}(F) = F_1 \cup \cdots \cup F_n$, then for each i, $\pi : F_i \to F$ is an isomorphism. Taking X to be $\pi^{-1}(U)$, the group G acts on X and $\pi : X \to U$ is a ramified cyclic covering of affine varieties. We consider the groups of units on the affine varieties U and X. In Section 3.1 these results are applied to the group of units on an affine curve.

Lemma 3.1. [7, Example II.6.5.1] *Let $F \subseteq \mathbb{P}^m$ be an irreducible hypersurface of degree d. For the open affine $U = \mathbb{P}^m - F$,*

(a) $\mathscr{O}^*(U)$, the group of units, is equal to k^*, and
(b) $\mathrm{Cl}(U)$, the class group, is a cyclic group of order d.

Proof. The degree homomorphism $\deg : \mathrm{Cl}(\mathbb{P}^m) \to \mathbb{Z}$ is an isomorphism. The exact sequence (1) becomes

$$1 \to k^* \to \mathscr{O}^*(U) \xrightarrow{\mathrm{div}} \mathbb{Z}F \xrightarrow{\chi} \mathrm{Cl}(\mathbb{P}^m) \to \mathrm{Cl}(U) \to 0 \tag{32}$$

where $\chi(F) = d$. Therefore, χ is one-to-one. □

Lemma 3.2. *In the context of Section 3.1, the following are true.*

(a) $(\mathscr{O}^*(X)/k^*)^G = \langle 1\rangle$.
(b) The $\mathbb{Z}$-module $\mathscr{O}^*(X)/k^*$ is free and has rank less than or equal to $n-1$.

Proof. Consider the commutative diagram

$$\begin{tikzcd}
1 \to \mathscr{O}^*(U)/k^* \arrow[r] \arrow[d, "\pi^\sharp"] & \mathbb{Z}F \arrow[r] \arrow[d, "\pi^*"] & \mathrm{Cl}(\mathbb{P}^m) \arrow[r] \arrow[d] & \mathrm{Cl}(U) \to 0 \arrow[d] \\
1 \to \mathscr{O}^*(X)/k^* \arrow[r, "\mathrm{div}"] & \bigoplus_{i=1}^{n} \mathbb{Z}F_i \arrow[r, "\chi"] & \mathrm{Cl}(Y) \arrow[r] & \mathrm{Cl}(X) \to 0
\end{tikzcd} \tag{33}$$

The top row of (33) is (32) and the second row is (1). The map π^* is defined by $F \mapsto F_1 + \cdots + F_n$. The maps on class groups are the natural maps. By [11, Lemma 10.34], the G-module $\bigoplus_{i=1}^n \mathbb{Z}F_i$ is free of rank one. Therefore,

$$\mathrm{H}^i\Big(G, \bigoplus_{i=1}^{n} \mathbb{Z}F_i\Big) = \begin{cases} \langle \pi^*(F) \rangle = \langle F_1 + \cdots + F_n \rangle & \text{if } i = 0 \\ (0) & \text{if } i > 0. \end{cases} \tag{34}$$

Let C denote the cokernel of π^*. Then C is a free $\mathbb{Z}$-module of rank $n-1$. The short exact sequence of G-modules $0 \to \mathbb{Z}F \xrightarrow{\pi^*} \bigoplus_i \mathbb{Z}F_i \to C \to 0$ gives the exact sequence

$$0 \to \mathbb{Z}F \xrightarrow{\pi^*} \langle \pi^*(F) \rangle \to C^G \xrightarrow{\partial^0} \mathrm{H}^1(G, \mathbb{Z}F) \to \cdots \tag{35}$$

of abelian groups. Since $\mathbb{Z}F$ is a trivial G-module, ∂^0 is the zero map and $C^G = (0)$. By Lemma 3.1, the cokernel of $\pi^\sharp$ is $\mathscr{O}^*(X)/k^*$. It follows from (33) that $\mathscr{O}^*(X)/k^*$ embeds as a G-submodule of C. This proves (a).

In the above context, if H is the subgroup of $\mathrm{Cl}(X)$ generated by the prime divisors $F_1, \ldots, F_p$, then (33) gives rise to the exact sequence

$$1 \to \mathscr{O}^*(X)/k^* \xrightarrow{\mathrm{div}} \bigoplus_{i=1}^{n} \mathbb{Z}F_i \xrightarrow{\chi} H \to 0 \tag{36}$$

of G-modules. A principal divisor on Y has degree 0. The subgroup of H generated by $\pi^*(F)$ is infinite cyclic. Part (b) follows from (36). □

Remark 3.3. The upper bound of Lemma 3.2(b) is sharp, as shown in Example 1.7.

Theorem 3.4. *In the above context, assume $n = p$ is a prime number. For the group $\mathscr{O}^*(X)/k^*$, there are two possibilities. The first is $\mathscr{O}^*(X) = k^*$, in which case H is a free $\mathbb{Z}$-module of rank p. The second possibility is that $\mathscr{O}^*(X)/k^*$ is a free $\mathbb{Z}$-module of rank $p-1$ and in this case H is isomorphic to an extension of $\mathbb{Z}$ by a finite group.*

Proof. As in Theorem 2.3, this follows from Lemma 3.2, the structure theory for a module over a cyclic group of order p, and (36). □

3.2. **The group of units on an affine curve.** In this section we apply Theorem 3.4 to study the group of units on an affine curve over $k = \mathbb{C}$, the field of complex numbers.

Let p be a prime number and $n \geq 3$ an integer such that $p \mid n$. Let $\lambda_1, \ldots, \lambda_n$ be distinct elements of k and set $f(x) = (x - \lambda_1) \cdots (x - \lambda_n)$. Let $X = Z(y^p - f(x))$, a nonsingular affine curve in $\mathbb{A}^2$. Let $\pi : Y \to \mathbb{A}^1$ be the morphism induced by $k[x] \to \mathscr{O}(X)$. Let Y be the complete nonsingular model for X and $\pi : Y \to \mathbb{P}^1$ the extension of π. Then Y is a cyclic cover of $\mathbb{P}^1$ of degree p, and we are in the context of the introduction to Section 3.

Let P_i denote the point on X (and on Y) where $y = x - \lambda_i = 0$. Let $Q_i = \pi(P_i)$. The map π ramifies at P_i and the ramification index is p. These are the only points where π is ramified. Solve the Riemann-Hurwitz Formula [7, Corollary IV.2.4] for the genus of Y to get $g(Y) = (p-1)(n-2)/2$.

The degree map $\deg : \mathrm{Cl}(Y) \to \mathbb{Z}$ is onto. The kernel of the degree map, denoted $\mathrm{Cl}^0(Y)$, is the Jacobian variety. By [9, p. 64], $\mathrm{Cl}^0(Y)$ is an abelian variety of dimension $g(Y)$. By [9, (iv), p. 42], $\mathrm{Cl}^0(Y)$ is a divisible group. Let P_0 be any closed point of Y. The group of divisors on Y of degree zero, denoted $\mathrm{Div}^0(Y)$, is generated by the set $\{P - P_0 \mid P \in \mathrm{Div}(Y)\}$. In this context, the exact sequence (1) becomes

$$1 \to \mathscr{O}^*(Y) \to \mathscr{O}^*(Y - P_0) \to \mathbb{Z}P_0 \xrightarrow{\chi} \mathrm{Cl}(Y) \to \mathrm{Cl}(Y - P_0) \to 0 \tag{37}$$

Since P_0 has degree one, the degree map is a splitting for χ, so $\mathrm{Cl}(Y - P_0)$ is isomorphic to $\mathrm{Cl}^0(Y)$.

Theorem 3.5. *Let $k = \mathbb{C}$ be the field of complex numbers and p a prime number. Say $G = \langle \sigma \rangle$ is a cyclic group of order p acting on a nonsingular projective curve Y such that the quotient map is $\pi : Y \to \mathbb{P}^1$. Assume Y has positive genus $g(Y) > 0$. For a sufficiently general $Q \in \mathbb{P}^1$, if $X = \pi^{-1}(\mathbb{P}^1 - Q)$, then $\mathscr{O}^*(X) = k^*$.*

Proof. Let $P_0 \in Y$ be a point where π is ramified. Let $Q_0 = \pi(P_0)$. For an arbitrary $P \in Y$, the divisor $\pi(P) - Q_0$ is a principal divisor on $\mathbb{P}^1$. Therefore,

$$\pi^*(\pi(P) - Q_0) = P + \sigma(P) + \cdots + \sigma^{p-1}(P) - pP_0 \tag{38}$$

is a principal divisor on Y. Manipulation of (38) shows that

$$(P - \sigma(P)) + 2\sigma(P - \sigma(P)) + 3\sigma^2(P - \sigma(P)) + \cdots + p\sigma^{p-1}(P - P_0) \tag{39}$$

is a principal divisor on Y. The group $\mathrm{Cl}^0(Y)$ is generated by the set of divisors $\{P - P_0 \mid P \in Y\}$ which is equal to the set $\{\sigma(P) - P_0 \mid P \in Y\}$. By (39) we see that $p\,\mathrm{Cl}^0(Y)$ is generated by the set of divisors $\{P - \sigma(P) \mid P \in Y\}$. The group $\mathrm{Cl}^0(Y)$ is an abelian variety of dimension $g(Y)$ over C. The torsion elements make up a discrete group which is isomorphic to $(\mathbb{Q}/\mathbb{Z})^{(2g)}$. For a sufficiently general choice of P, $P - \sigma(P)$ generates an infinite subgroup of $\mathrm{Cl}^0(Y)$. Let H denote the subgroup of $\mathrm{Cl}(Y)$ generated by $P, \sigma(P), \ldots, \sigma^{p-1}(P)$. Then H contains the infinite subgroup $\langle P - \sigma(P), \sigma(P) - \sigma^2(P), \ldots, \sigma^{p-1}(P) - P \rangle$. If $Q = \pi(P)$ and $X = Y - \pi^{-1}(Q) = \pi^{-1}(\mathbb{P}^1 - Q)$, then Theorem 3.4 implies X has a trivial group of units, $\mathscr{O}^*(X) = k^*$. □

Proposition 3.6. *Let $k = \mathbb{C}$ be the field of complex numbers. Let p be a prime number, $n \geq 3$ an integer such that $p \mid n$, and $\lambda_1, \ldots, \lambda_n$ distinct elements in k. Let $f(x) = \prod_{i=1}^n (x - \lambda_i)$ and $X = Z(y^p - f(x))$, an affine curve in $\mathbb{A}^2$. For a sufficiently general choice of $\lambda_1, \ldots, \lambda_n$, the group of units on X is equal to k^*.*

Proof. Apply Theorem 3.5 to the nonsingular projective completion of X. □

Proposition 3.7. *Let $k = \mathbb{C}$, the field of complex numbers. Let $n \geq 3$ and $\lambda_1, \ldots, \lambda_n$ distinct elements in k. Let $f(x) = \prod_{i=1}^n (x - \lambda_i)$ and $X = Z(y^2 - f(x))$ the corresponding affine hyperelliptic curve in $\mathbb{A}^2$. For a sufficiently general choice of $\lambda_1, \ldots, \lambda_n$, the group of units on X is equal to k^*.*

Proof. Let Y be the nonsingular projective completion for X. If n is even, this follows from Theorem 3.5. If n is odd, then $Y - X$ consists of only one point and we apply (37). □

Example 3.8. In Proposition 3.7, if X is arbitrary, the group of units can be non-trivial. For instance, let $f = x^4 - 1$. In $\mathscr{O}(X)$, we have $1 = x^4 - y^2$, hence $x^2 - y$ is invertible. For another example, let $f = x^4 + x$ and $Y = Z(y^2 - f)$. In $\mathscr{O}(Y)$,

$$\begin{aligned}\big((x^3+1/2)+xy\big)\big((x^3+1/2)-xy\big) &= (x^3+1/2)^2 - x^2y^2 \\ &= x^6 + x^3 + 1/4 - x^2(x^4+x) \\ &= 1/4\end{aligned}$$

Hence $\mathscr{O}^*(X) \neq k^*$.

3.3. **The ramification divisor is at infinity.** Start with a normal projective variety Y of dimension $m > 0$, together with $G = \langle \sigma \rangle$ a cyclic group of order n acting on Y such that the quotient morphism is $\pi : Y \to \mathbb{P}^m$. Assume the ramification divisor of π on Y is the set of prime divisors $Q = \{Q_1, \dots, Q_r\}$ and that the ramification index at each Q_i is n. Let $\pi(Q) = P = \{P_1, \dots, P_r\}$. Let $U = \mathbb{P}^m - P$ and $X = Y - Q$. Then $\pi : X \to U$ is a Galois cover with group G. The Nagata sequences for U and X give the rows of the commutative diagram

$$\begin{tikzcd}
1 \to \mathscr{O}^*(U)/k^* \arrow[r] \arrow[d, "\pi^\sharp"'] & \bigoplus_{i=1}^{r} \mathbb{Z}P_i \arrow[r] \arrow[d, "\delta"] & \mathrm{Cl}(\mathbb{P}^m) \arrow[r] \arrow[d] & \mathrm{Cl}(U) \to 0 \arrow[d] \\
1 \to \mathscr{O}^*(X)/k^* \arrow[r] & \bigoplus_{i=1}^{r} \mathbb{Z}Q_i \arrow[r, "\chi"] & \mathrm{Cl}(Y) \arrow[r] & \mathrm{Cl}(X) \to 0
\end{tikzcd} \tag{40}$$

By the map δ, P_i is mapped to nQ_i [6, p. 30]. We consider the group of units $\mathscr{O}^*(X)/k^*$ and the image of χ, which is equal to the subgroup of $\mathrm{Cl}(Y)$ generated by $Q_1, \dots, Q_r$.

In this section we utilize étale cohomology. The étale cohomology group $\mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z})$ classifies the Galois covers of U with finite cyclic group.

Proposition 3.9. *In the above context, if* $\mathrm{Cl}(U) = \langle 0 \rangle$*, then*
(a) $\mathscr{O}^*(U)/k^*$ *is a free* $\mathbb{Z}$*-module of rank* $r-1$*,*
(b) $\mathscr{O}^*(X)/k^*$ *is a free* $\mathbb{Z}$*-module of rank* $r-1$*, and*
(c) $\mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z})$ *is a free* $\mathbb{Q}/\mathbb{Z}$*-module of rank* $r-1$*.*

Proof. For each $\nu > 1$ the diagram

$$\begin{tikzcd}
1 \arrow[r] & \mathscr{O}^*(U)/k^* \arrow[r] \arrow[d, "\nu"] & \bigoplus_{i=1}^{r} \mathbb{Z}P_i \arrow[r] \arrow[d, "\nu"] & \mathrm{Cl}(\mathbb{P}^m) \arrow[r] \arrow[d, "\nu"] & 0 \\
1 \arrow[r] & \mathscr{O}^*(U)/k^* \arrow[r] \arrow[d] & \bigoplus_{i=1}^{r} \mathbb{Z}P_i \arrow[r] \arrow[d] & \mathrm{Cl}(\mathbb{P}^m) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathrm{H}^1(U, \mu_\nu) \arrow[r] & (\mathbb{Z}/\nu)^{(r)} \arrow[r] & \mathbb{Z}/\nu \arrow[r] & 0
\end{tikzcd} \tag{41}$$

commutes. The first and second rows are from (40), and are split exact since $\mathrm{Pic}(\mathbb{P}^m) \cong \mathbb{Z}$ and $\mathrm{Pic}(U) = \langle 0 \rangle$. This gives (a). In (40) the image of χ is infinite and $\pi^\sharp$ is one-to-one, which implies (b). The vertical maps connecting the first and second rows of (41) are "multiplication by ν" and are one-to-one. The first column of (41) is the Kummer sequence. The vertical maps connecting the second and third rows are onto, since $\operatorname{Pic}(U) = \langle 0 \rangle$. The third row is split exact. To get (c), take the limit over all ν. $\square$

Proposition 3.10. *Let $\pi\colon X \to U$ be a cyclic Galois cover of degree n of integral varieties with group G. The sequence*

$$0 \to \mathbb{Z}/n \to \mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z}) \xrightarrow{\pi^*} \mathrm{H}^1(X, \mathbb{Q}/\mathbb{Z})^G \to 0$$

is exact. The kernel of π^* is generated by the class represented by the cyclic covering $\pi : X \to U$.

Proof. The Hochschild-Serre spectral sequence [8, p. 105] for $\pi : X \to U$ is

$$\mathrm{H}^p(G, \mathrm{H}^q(X, \mathbb{Q}/\mathbb{Z})) \Rightarrow \mathrm{H}^{p+q}(U, \mathbb{Q}/\mathbb{Z}). \tag{42}$$

Since X is integral, the group of global sections of the constant sheaf is $\mathrm{H}^0(X, \mathbb{Q}/\mathbb{Z}) = \mathbb{Q}/\mathbb{Z}$. Because G acts trivially on $\mathbb{Q}/\mathbb{Z}$, $\mathrm{H}^0(X, \mathbb{Q}/\mathbb{Z})^G = \mathbb{Q}/\mathbb{Z}$. By [11, Theorem:10.35]

$$\mathrm{H}^p(G, \mathbb{Q}/\mathbb{Z}) = \begin{cases} 0 & \text{if } p > 0 \text{ is even} \\ \mathbb{Z}/n & \text{if } p \text{ is odd.} \end{cases}$$

Consider the filtration $\mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z}) = E_0^1 \supseteq E_1^1 \supseteq 0$. Then

$$E_1^1 = E_\infty^{1,0} = E_2^{1,0} = \mathrm{H}^1(G, H^0(X, \mathbb{Q}/\mathbb{Z})) \cong \mathbb{Z}/n$$

and since $d_2^{0,1} : \mathrm{H}^1(X, \mathbb{Q}/\mathbb{Z})^G \to \mathrm{H}^2(G, \mathbb{Q}/\mathbb{Z})$ is the zero map, $E_0^1/E_1^1 = E_\infty^{0,1} = E_2^{0,1} = \mathrm{H}^1(X, \mathbb{Q}/\mathbb{Z})^G$. Therefore we have the short exact sequence

$$0 \to \mathbb{Z}/n \to \mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z}) \to \mathrm{H}^1(X, \mathbb{Q}/\mathbb{Z})^G \to 0.$$

The inclusion $E_1^1 \subseteq E_0^1$ corresponds to the subgroup of $\mathrm{H}^1(U, \mathbb{Q}/\mathbb{Z})$ generated by the cyclic Galois cover $\pi : X \to U$. $\square$

Lemma 3.11. *In the context of Section 3.3, the following are true.*

(a) The image of $\pi^* : \mathrm{H}^1(U, \mu_n) \to \mathrm{H}^1(X, \mu_n)$ is a subgroup of the image of $\rho : \mathrm{H}^1(Y, \mu_n) \to \mathrm{H}^1(X, \mu_n)$. We say that π^* splits the ramification of each $\xi \in \mathrm{H}^1(U, \mu_n)$.

(b) If $r \geq 2$ and $\operatorname{Cl}(U) = \langle 0 \rangle$, the image of π^* is isomorphic to $(\mathbb{Z}/n)^{(r-2)}$.

Proof. By $\operatorname{Sing} Y$ we denote the singular locus of Y. Since Y is normal, the codimension of $\operatorname{Sing} Y$ in Y is at least two. The codimension of $\operatorname{Sing} Q$ in Y is at least two. Since $X \to U$ is unramified, we know X is nonsingular. Let ω be a closed subset of Q, of codimension greater than or equal to two, such that $Y - \omega$ and $Q - \omega$ are nonsingular. Because π is a finite morphism, $\pi(\omega)$ has codimension at least two in $\mathbb{P}^m$. By [8, Lemma VI.9.1], $\mathrm{H}^1(Y, \mu_n) = \mathrm{H}^1(Y - \omega, \mu_n)$. The diagram

$$\begin{tikzcd}
\mathrm{H}^1_{P_i - \pi(\omega)}(\mathbb{P}^m - \pi(\omega), \mathbb{G}_m) \arrow[r] \arrow[d, "\alpha"] & \mathrm{H}^1(\mathbb{P}^m - \pi(\omega), \mathbb{G}_m) \arrow[d, "\pi^*"] \\
\mathrm{H}^1_{Q_i - \omega}(Y - \omega, \mathbb{G}_m) \arrow[r] & \mathrm{H}^1(Y - \omega, \mathbb{G}_m)
\end{tikzcd} \tag{43}$$

commutes and α agrees with the map δ of diagram (40). That is, α is "multiplication by n". Consider the commutative diagram

$$\begin{tikzcd}
\mathrm{H}^1_{P_i-\pi(\omega)}(\mathbb{P}^m-\pi(\omega),\mathbb{G}_m) \arrow[r] \arrow[d, "\alpha"] & \mathrm{H}^2_{P_i-\pi(\omega)}(\mathbb{P}^m-\pi(\omega),\mu_n) \arrow[d, "\beta"] \\
\mathrm{H}^1_{Q_i-\omega}(Y-\omega,\mathbb{G}_m) \arrow[r] & \mathrm{H}^2_{Q_i-\omega}(Y-\omega,\mu_n)
\end{tikzcd} \tag{44}$$

Then on the class of $P_i-\pi(\omega)$ in $\mathrm{H}^2_{P_i-\pi(\omega)}(\mathbb{P}^m-\pi(\omega),\mu_n)$, β is "multiplication by n". Consider the commutative diagram

$$\begin{tikzcd}
\mathrm{H}^1(\mathbb{P}^m-\pi(\omega),\mu_n) \arrow[r] \arrow[d] & \mathrm{H}^1(U,\mu_n) \arrow[r, "\partial^0"] \arrow[d, "\pi^*"] & \bigoplus_{i=1}^{r}\mathrm{H}^2_{P_i-\pi(\omega)}(\mathbb{P}^m-\pi(\omega),\mu_n) \arrow[d, "\gamma"] \\
\mathrm{H}^1(Y-\omega,\mu_n) \arrow[r, "\rho"] & \mathrm{H}^1(X,\mu_n) \arrow[r] & \bigoplus_{i=1}^{r}\mathrm{H}^2_{Q_i-\omega}(Y-\omega,\mu_n)
\end{tikzcd} \tag{45}$$

On the image of ∂^0, γ is "multiplication by n". The image of π^* is a subgroup of the image of ρ. In part (b), Proposition 3.10 implies that the kernel of π^* in (45) is cyclic of order n. By Proposition 3.9, the image of π^* in (45) is isomorphic to $(\mathbb{Z}/n)^{(r-2)}$. $\square$

Proposition 3.12. *In the notation of Section 3.3, assume* $\mathrm{Cl}(U)=\langle 0\rangle$ *and* $r\geq 2$.

- *(a) The Galois cover $X\to U$ corresponds to adjoining the nth root of an element $f\in\mathscr{O}^*(U)$. That is, $\mathscr{O}(X)$ is isomorphic to $\mathscr{O}(U)[z]/(z^n-f)$.*
- *(b) Let H denote the image of χ in* (40), *which is the subgroup of* $\mathrm{Cl}(Y)$ *generated by $Q_1,\dots,Q_r$.*
  - *(a) The $\mathbb{Z}/n$-rank of $H\otimes\mathbb{Z}/n$ is equal to $r-1$.*
  - *(b) The $\mathbb{Z}/n$-rank of ${}_nH$ is equal to $r-2$.*
- *(c) If $f_1,\dots,f_{r-1}$ is a basis for $\mathscr{O}^*(U)/k^*$, then the group $\mathscr{O}^*(X)/k^*$ is generated by $z,f_1,\dots,f_{r-1}$.*

Proof. There is a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathscr{O}^*(U)\otimes\mathbb{Z}/n \arrow[r, "="] \arrow[d, "\pi^\sharp"] & \mathrm{H}^1(U,\mathbb{Z}/n) \arrow[r] \arrow[d, "\pi^*"] & 0 \\
0 \arrow[r] & \mathscr{O}^*(X)\otimes\mathbb{Z}/n \arrow[r] & \mathrm{H}^1(X,\mathbb{Z}/n) \arrow[r] & {}_n\mathrm{Cl}(X)
\end{tikzcd} \tag{46}$$

so by Proposition 3.10, the kernel of $\pi^\sharp$ is cyclic of order n. This is (a). From (40) we have the commutative diagram

$$\begin{tikzcd}
1 \arrow[r] & \mathscr{O}^*(X)/k^* \arrow[r] \arrow[d, "n"] & \bigoplus_{i=1}^r \mathbb{Z}Q_i \arrow[r, "\chi"] \arrow[d, "n"] & H \arrow[r] \arrow[d, "n"] & 0 \\
1 \arrow[r] & \mathscr{O}^*(X)/k^* \arrow[r] & \bigoplus_{i=1}^r \mathbb{Z}Q_i \arrow[r, "\chi"] & H \arrow[r] & 0
\end{tikzcd} \tag{47}$$

where the vertical maps are "multiplication by n". The Snake Lemma gives the exact sequence

$$0 \to {}_nH \to (\mathbb{Z}/n)^{(r-1)} \to (\mathbb{Z}/n)^{(r)} \to H \otimes \mathbb{Z}/n \to 0 \tag{48}$$

Let $f_1, \dots, f_{r-1}$ be a basis for $\mathscr{O}^*(U)/k^*$. Then $\operatorname{div}\left(\pi^\sharp(f_i)\right)$ is divisible by n, hence H is a homomorphic image of $\mathbb{Z} \oplus (\mathbb{Z}/n)^{(r-1)}$. But $\operatorname{div}(z)$ is not divisible by n. Since X is integral, we know that $\operatorname{div}(z)$ is a non-trivial relation. Therefore, H is a homomorphic image of $\mathbb{Z} \oplus (\mathbb{Z}/n)^{(r-2)}$. The group ${}_nH$ of elements annihilated by n contains the image of π^* which by Lemma 3.11 has rank equal to $r-2$. ☐

In the notation of Section 3.3, Proposition 3.13 is the $r = 1$ case.

Proposition 3.13. *Let P be an irreducible divisor on $\mathbb{P}^m$ of degree d and $U = \mathbb{P}^m - P$. Let $X \to U$ be a cyclic Galois cover of degree d corresponding to a generator of $\mathrm{H}^1(U, \mu_d)$. Then $\mathscr{O}^*(X) = k^*$.*

Proof. The Galois cover X exists by Lemma 3.1. Let $\pi : Y \to \mathbb{P}^m$ be the cyclic cover which ramifies along P. Let $\pi^{-1}(P) = Q$. In diagram (40), $\chi(Q)$ is non-zero because a principal divisor on Y has degree zero [7, Exercise II.6.2]. ☐

3.4. **The group of units on an affine variety.** In this section we apply results from Section 3.3 to derive the group of units on the affine variety defined by an equation of the form $f_1 \cdots f_n = 1$, where the polynomials f_i are distinct linear forms in $k[x_1, \dots, x_m]$.

Example 3.14. This is a special case of Example 1.4. Let $m \geq 1$ and $n \geq 2$. Let $f_1, \dots, f_n$ be linear forms in $k[x_1, \dots, x_m]$ and Y the projective variety in $\mathbb{P}^m$ defined by $f_1 \cdots f_n = x_0^n$. Let $Z_0 = Z(x_0)$ denote the hyperplane at infinity. For each $i = 1, \dots, n$, let $F_i = Z(f_i)$ be the hyperplane in $\mathbb{P}^m$ defined by f_i and set $L_i = F_i \cap Z_0$. The divisor at infinity on Y is $L_1 + \cdots + L_n$. View Y as a cyclic cover of $Z_0 = \operatorname{Proj} k[x_1, \dots, x_m]$ and let $\pi : Y \to Z_0$ be the projection. The ramification divisor of π is the set $L = \{L_1, \dots, L_n\}$. Let X be the open affine $Y - L$. Then $\mathscr{O}(X) = k[x_1, \dots, x_m]/(f_1 \cdots f_n - 1)$. In $\mathscr{O}(X)$ the functions f_i are invertible. One can compute the divisors $\operatorname{div}(f_i) = nL_i - L_1 - \cdots - L_n$ on Y. Consider the subgroup of $\bigoplus_{i=1}^n \mathbb{Z}L_i$ generated by $\operatorname{div}(f_1), \dots, \operatorname{div}(f_n)$. The invariant factors of the matrix

$$\begin{bmatrix}
n-1 & -1 & \dots & -1 & -1 \\
-1 & n-1 & \dots & -1 & -1 \\
 & & \vdots & & \\
-1 & -1 & \dots & n-1 & -1 \\
-1 & -1 & \dots & -1 & n-1
\end{bmatrix} \tag{49}$$

are 1 and n, where n occurs with multiplicity $n-2$. By Proposition 3.12 we know that H is isomorphic to $\mathbb{Z} \oplus (\mathbb{Z}/n)^{(n-2)}$. Therefore, $\{f_1, \dots, f_{n-1}\}$ is a basis for the group $\mathscr{O}^*(X)/k^*$.

REFERENCES

[1] S. U. Chase, D. K. Harrison, and Alex Rosenberg, *Galois theory and Galois cohomology of commutative rings*, Mem. Amer. Math. Soc. No. **52** (1965), 15–33. MR 0195922 (33 #4118)

[2] Charles W. Curtis and Irving Reiner, *Representation theory of finite groups and associative algebras*, Pure and Applied Mathematics, Vol. XI, Interscience Publishers, a division of John Wiley & Sons, New York-London, 1962. MR 0144979 (26 #2519)

[3] T. J. Ford, *The Brauer group of an affine cone*, J. Pure Appl. Algebra **155** (2001), no. 1, 29–40. MR 1804326 (2001k:14044)

[4] Timothy J. Ford, *The relative Brauer group of a cyclic cover of affine space*, J. Pure Appl. Algebra **215** (2011), no. 5, 847–865. MR 2747225 (2012a:16036)

[5] ______, *The relative Brauer group of an affine double plane*, Comm. Algebra (2012), to appear.

[6] Robert M. Fossum, *The divisor class group of a Krull domain*, Springer-Verlag, New York, 1973. MR 0382254 (52 #3139)

[7] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977. MR 0463157 (57 #3116)

[8] James S. Milne, *Étale cohomology*, Princeton University Press, Princeton, N.J., 1980. MR 559531 (81j:14002)

[9] David Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, No. 5, Published for the Tata Institute of Fundamental Research, Bombay, 1970. MR 0282985 (44 #219)

[10] David E. Rohrlich, *Points at infinity on the Fermat curves*, Invent. Math. **39** (1977), no. 2, 95–127. MR 0441978 (56 #367)

[11] Joseph J. Rotman, *An introduction to homological algebra*, Pure and Applied Mathematics, vol. 85, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1979. MR 538169 (80k:18001)

[12] Pierre Samuel, *À propos du théorème des unités*, Bull. Sci. Math. (2) **90** (1966), 89–96. MR 0204454 (34 #4296)

[13] Jacques Vélu, *Sur un théorème de D. Rohrlich*, C. R. Acad. Sci. Paris Sér. A-B **290** (1980), no. 3, A157–A160. MR 563965 (81f:14013)

E-mail address: `ford@fau.edu`

DEPARTMENT OF MATHEMATICS, FLORIDA ATLANTIC UNIVERSITY, BOCA RATON, FLORIDA 33431